



## Missouri Pharmacy Program – Preferred Drug List



### Tramadol-like Agents

*Effective 06/21/2012*

*Revised 04/07/2016*

#### Preferred Agents

- Tramadol
- Tramadol-Apap

#### Non-Preferred Agents

- Conzip®
- Nucynta®
- NucyntaER®
- Rybix® ODT
- Ryzolt ER®/ODT
- Tramadol ER
- Ultracet®
- Ultram®
- Ultram ER®

<u>Approval Criteria</u>	<u>Denial Criteria</u>
Failure to achieve desired therapeutic outcomes with trial on 2 or more preferred agents <ul style="list-style-type: none"> <li>• Documented trial period for preferred agents</li> <li>• Documented ADE/ADR to preferred agents</li> </ul>	(see below)
Documented compliance on current therapy regimen	Drug Prior Authorization Hotline: (800) 392-8030

### Denial Criteria

#### **Ultram:**

- 15 years of age or less<sup>2</sup>
- Daily dose exceeds maximum recommended daily dose:
  - Patients ≤ 75 years of age – 400 mg
  - Patients > 75 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

#### **Ultracet:**

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients ≥ 16 years of age – 300 mg tramadol and 2600mg acetaminophen

- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

#### **Ultram ER/Ryzolt:**

- 15 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  16 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days
- History of opioid abuse or dependence – therapy will be subject to Clinical Consultant review
- Claim for Ultram ER/Ryzolt therapy without previous stabilization on immediate-release tramadol product.

#### **Nucynta:**

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  18 years of age – 700 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

#### **Nucynta ER:**

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  18 years of age – 500 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

#### **Rybix ODT**

- 16 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  17 years of age – 400mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days