

**NEW DRUG REVIEWS FOR 12/17/2009 PA MEETING**

**OPEN ACCESS**

**FOR INFORMATION ONLY**

**COMMON TRADE NAME**

**GENERIC NAME**

**INDICATIONS**

| COMMON TRADE NAME                                  | GENERIC NAME                   | INDICATIONS  |
|--|--------------------------------|--|
| Aplenzin ER (Tablet 174 mg)                        | Bupropion HBR                  | Indicated for the treatment of depression in adults.   |
| Caldolor (Vial 400 mg/4 ml)                        | Ibuprofen                      | Indicated to treat pain and fever.   |
| Caldolor (Vial 800 mg/8 ml)                        | Ibuprofen                      | Indicated to treat pain and fever.   |
| Creon (Capsule DR 120K-24K-76K)                    | Amylase/Lipase/Protease        | Indicated for the treatment of exocrine pancreatic insufficiency due to cystic fibrosis or other conditions.   |
| Creon (Capsule DR 30K-6K19K)                       | Amylase/Lipase/Protease        | Indicated for the treatment of exocrine pancreatic insufficiency due to cystic fibrosis or other conditions.   |
| Creon (Capsule DR 60K-12K-38K)                     | Amylase/Lipase/Protease        | Indicated for the treatment of exocrine pancreatic insufficiency due to cystic fibrosis or other conditions.   |
| Gesticare DHA (CMB Tablet MP 27-1-250 mg)          | PNV w CA, NO.34/iron/FA/DHA    | Prenatal Vitamin   |
| Hectorol (Capsule 1 mcg)                           | Doxercalciferol                | Indicated for the treatment of secondary hyperparathyroidism in patients with Stage 3 or Stage 4 Chronic Kidney Disease (capsules) and in patients with Chronic Kidney Disease on dialysis (capsules and injection). |
| IC400 (Combo Package 400 mg -65 mg)                | Ibuprofen/Caff/B1/B2/B6/B12    | Used for treatment of mild to moderate pain  |
| IC800 (Combo Package 800 mg-65 mg)                 | Ibuprofen/Caff/B1/B2/B6/B12    | Used for treatment of mild to moderate pain  |
| Invega Sustenna (Disposable Syringe 117 mg/0.75)   | Paliperidone Palmitate         | Used for the treatment of schizophrenia.   |
| Invega Sustenna (Disposable Syringe 156 mg/1 ml)   | Paliperidone Palmitate         | Used for the treatment of schizophrenia.   |
| Invega Sustenna (Disposable Syringe 234 mg/1.5 ml) | Paliperidone Palmitate         | Used for the treatment of schizophrenia.   |
| Invega Sustenna (Disposable Syringe 39 mg-0.25)    | Paliperidone Palmitate         | Used for the treatment of schizophrenia.   |
| Invega Sustenna (Disposable Syringe 78 mg/0.5)     | Paliperidone Palmitate         | Used for the treatment of schizophrenia.   |
| Ivites Rx (Tablet 1 mg-60 mg)                      | VIT B CMPLX NO3/FA/C/BIOT/ZINC | Multivitamin product used to treat or prevent vitamin deficiency due to poor diet, certain illnesses, or during pregnancy.   |
| Ilaris (Vital 180 mg)                              |                                | Used in the treatment of CADS  |
| Multaq (Tablet 400 mg)                             | Dronedarone Hydrochloride      | Indicated to help maintain normal heart rhythms in patients with a history of atrial fibrillation or atrial flutter (heart rhythm disorders).  |
| Nucynta (Talbet 100 mg)                            | Tapentadol Hydrochloride       | Used to manage moderate to severe pain. It works by acting in the brain to relieve pain.   |
| Nucynta (Talbet 50 mg)                             | Tapentadol Hydrochloride       | Used to manage moderate to severe pain. It works by acting in the brain to relieve pain.   |
| Nucynta (Talbet 75 mg)                             | Tapentadol Hydrochloride       | Used to manage moderate to severe pain. It works by acting in the brain to relieve pain.   |
| Ozurdex (Implant 0.7 mg)                           | Dexamethasone                  | Indicated for the treatment of macular edema following branch retinal vein occlusion (BRVO) or central retinal vein occlusion (CRVO).  |
| Plan B One-Step (Tablet 1.5 mg)                    | Levonorgestrel                 | Intended to prevent pregnancy after known or suspected contraceptive failure or unprotected intercourse.   |
| Prefera-OB Plus DHA (Combo Package 22-6-1-200)     | PNV COMBO #8/iron/Folic AC/DHA | Prenatal Vitamin   |
| Saphris (Tablet sublingual 10 mg)                  | Asenapine                      | indicated for acute treatment of schizophrenia in adults. Acute treatment of manic or mixed episodes associated with bipolar I disorder in adults.   |
| Saphris (Tablet sublingual 5 mg)                   | Asenapine                      | indicated for acute treatment of schizophrenia in adults. Acute treatment of manic or mixed episodes associated with bipolar I disorder in adults.   |
| Tyvaso (1.7 mg/2.9 ml Ampul)                       |                                | Used in the treatment of Pulmonary Hypertension  |

|                                |          |   |
|--------------------------------|----------|---|
| Ventavis (Solution 20 mcg/1ml) | Iloprost | Indicated for the treatment of pulmonary arterial hypertension (WHO Group I) in patients with NYHA Class III or IV symptoms. In controlled trials, it improved a composite endpoint consisting of exercise tolerance, symptoms (NYHA Class), and lack of deterioration. |
|--------------------------------|----------|---|

**PREFERRED DRUG LIST IMPLEMENTATION**  
**COMMON TRADE NAME**

**GENERIC NAME**

**INDICATIONS**

|   |                                |  |
|---|--------------------------------|--|
| Acuvail (Droperette 0.45%)                      | Ketorolac Tromethamine         | Ophthalmic NSAIDS <b>NON PREFERRED</b>   |
| Adcirca (Tablet 20 mg)                          | Tadalafil                      | Oral Pulmonary Hypertension PDL Edit <b>NON PREFERRED</b>  |
| Benzaclin (kit 1%-5%)                           | Clindamycin/Benz Per/Hyalur NA | Benzoyl Peroxide/Clindamycin Combo PDL Edit <b>PREFERRED</b>   |
| Bepreve (Drops 1.5%)                            | Bepotastine Besilate           | Ophthalmic Antihistamine PDL Edit <b>NON PREFERRED</b>   |
| Edular (Tablet Sublingual 10 mg)                | Zolpidem Tartrate              | Indicated for the short-term treatment of insomnia characterized by difficulties with sleep initiation. <b>NON PREFERRED</b>   |
| Edular (Tablet Sublingual 5 mg)                 | Zolpidem Tartrate              | Indicated for the short-term treatment of insomnia characterized by difficulties with sleep initiation. <b>NON PREFERRED</b>   |
| Effient (Tablet 10 mg)                          | Prasurgel Hydrochloride        | Indicated to reduce the rate of thrombotic cardiovascular (CV) events (including stent thrombosis) in patients with acute coronary syndrome (ACS) who are to be managed with percutaneous coronary intervention (PCI) as follows:<br><ul style="list-style-type: none"> <li>• Patients with unstable angina (UA) or non-ST-elevation myocardial infarction (NSTEMI).</li> <li>• Patients with ST-elevation myocardial infarction (STEMI) when managed with primary or delayed PCI. <b>NON PREFERRED</b></li> </ul> |
| Effient (Tablet 5 mg)                           | Prasurgel Hydrochloride        | Indicated to reduce the rate of thrombotic cardiovascular (CV) events (including stent thrombosis) in patients with acute coronary syndrome (ACS) who are to be managed with percutaneous coronary intervention (PCI) as follows:<br><ul style="list-style-type: none"> <li>• Patients with unstable angina (UA) or non-ST-elevation myocardial infarction (NSTEMI).</li> <li>• Patients with ST-elevation myocardial infarction (STEMI) when managed with primary or delayed PCI. <b>NON PREFERRED</b></li> </ul> |
| Embeda (Capsule SR Pel 100 mg-4 mg)             | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Embeda (Capsule SR Pel 20 mg-0.8 mg)            | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Embeda (Capsule SR Pel 30 mg-1.2 mg)            | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Embeda (Capsule SR Pel 50 mg-2 mg)              | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Embeda (Capsule SR Pel 60 mg-2.4 mg)            | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Embeda (Capsule SR Pel 80 mg-3.2 mg)            | Morphine Sulfate/Naltrexone    | Long Acting Narcotics <b>NON PREFERRED</b>   |
| Fibricor (Tablet 105 mg)                        | Fenofibric Acid                | Triglyceride Lowering Agents <b>NON PREFERRED</b>  |
| Fibricor (Tablet 35 mg)                         | Fenofibric Acid                | Triglyceride Lowering Agents <b>NON PREFERRED</b>  |
| Norditropin Nordiflex (Pen Injector 30 mg/3 ml) | Somatropin                     | Human Growth Hormone <b>PREFERRED</b>  |
| Onglyza (Tablet 2.5 mg)                         | Saxagliptin Hydrochloride      | DPP-IV PDL Edit <b>NON PREFERRED</b>   |
| Onglyza (Tablet 5 mg)                           | Saxagliptin Hydrochloride      | DPP-IV PDL Edit <b>NON PREFERRED</b>   |
| Renvela (Powder Packet 0.8 G)                   | Sevelamer Carbonate            | Electrolyte Depleters <b>NON PREFERRED</b>   |
| Renvela (Powder Packet 2.4 G)                   | Sevelamer Carbonate            | Electrolyte Depleters <b>NON PREFERRED</b>   |
|   |                                |  |

**CLINICAL EDITS FOR IMPLEMENTATION****COMMON TRADE NAME****GENERIC NAME****INDICATIONS**

|                              |                      |   |
|------------------------------|----------------------|---|
| Intuniv (Tablet SR 24H 1 mg) | Guanfacine HCl       | Indicated for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children and adolescents aged 6 to 17 years |
| Intuniv (Tablet SR 24H 2 mg) | Guanfacine HCl       | Indicated for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children and adolescents aged 6 to 17 years |
| Intuniv (Tablet SR 24H 3 mg) | Guanfacine HCl       | Indicated for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children and adolescents aged 6 to 17 years |
| Intuniv (Tablet SR 24H 4 mg) | Guanfacine HCl       | Indicated for the treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in children and adolescents aged 6 to 17 years |
| Sabril (Tablet 500 mg)       | Vigabatrin           | 15 Day Limitation Clinical Edit   |
| Sarafem (Tablet 15mg)        | Fluoxetine HCL       | Selective Serotonin Reuptake Inhibitor (SSRI)   |
| Zipsor (Capsule 25 mg)       | Diclofenac Potassium | NSAID Step Therapy Edit   |
|                              |                      |   |
|                              |                      |   |

**REQUEST FOR CONTINUED PRIOR AUTHORIZATION****COMMON TRADE NAME****GENERIC NAME****INDICATIONS**

|                                   |                                |   |
|-----------------------------------|--------------------------------|---|
| Analpram E (Kit 2.5-1% (4G))      | Hydrocor/Pram/Emol/Skn clns#18 | Indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.   |
| Aloquin (Gel(GM) 1.25%-1%)        | Iodoquinol/Aloe Polysacchar#1  | "Possibly" Effective: Contact or atopic dermatitis; impetiginized eczema; nummular eczema; endogenous chronic infectious dermatitis; stasis dermatitis; pyoderma; nuchal eczema and chronic eczematoid otitis externa; acne urticata; localized or disseminated neurodermatitis; lichen simplex chronicus; anogenital pruritus (vulvae, scroti, ani); folliculitis; bacterial dermatoses; mycotic dermatoses such as tinea (capitis, cruris, corporis, pedis); monliasis; intertrigo. |
| Benzefoam (Foam 5.3%)             | Benzoyl Peroxide               | Indicated for use in the topical treatment of mild to moderate acne vulgaris.   |
| Notuss-PE (Liquid 10-10 mg/5 ml)  | Phenylephrine HCL/Codeine PHOS | Cough and Cold  |
| Ryneze (Liquid 4-1.25 mg/5 ml)    | Chlor-Mal/Methscopolamine NIT  | Cough and Cold  |
| Solodyn (Tablet SR 24H 115 mg)    | Minocycline HCl                | An oral antibiotic approved to treat only the red, pus-filled pimples and red bumps of moderate to severe acne in patients 12 years of age and older.   |
| Solodyn (Tablet SR 24H 65 mg)     | Minocycline HCl                | An oral antibiotic approved to treat only the red, pus-filled pimples and red bumps of moderate to severe acne in patients 12 years of age and older.   |
| Tekral (Tablet SR 12H 120-100 mg) | P-Ephed HCL/DP-Hydrum HCl      | Cough and Cold  |
| Triaz (Foaming Cloths 3%)         | Benzoyl Peroxide               | Indicated for the treatment of acne.  |
| Uramaxin (Lotion 45%)             | Urea                           | Used to treat dry/rough skin conditions (e.g., eczema, psoriasis, corns, callus) and some nail problems (e.g., ingrown nails). It may also be used to help remove dead tissue in some wounds to help wound healing.   |
| Zotex-Ex (Tablet 350-15-12 mg)    | Guaifenesin/D-Methorphan HB/PE | Cough and Cold  |