

SmartPA Criteria Proposal

Drug/Drug Class:	Antihyperuricemic Agents PDL Edit
First Implementation Date:	June 21, 2011
Proposed Date:	September 15, 2022
Prepared For:	MO HealthNet
Prepared By:	MO HealthNet/Conduent
Criteria Status:	<input checked="" type="checkbox"/> Existing Criteria <input type="checkbox"/> Revision of Existing Criteria <input type="checkbox"/> New Criteria

Executive Summary

Purpose: The MO HealthNet Pharmacy Program will implement a state-specific preferred drug list.

Why Issue Selected: Hyperuricemia, defined as serum uric acid greater than 6.8 mg/dL, can occur either due to an overproduction of uric acid, an under excretion of uric acid, or a combination of the two mechanisms. Most often, hyperuricemia results as a reduction in fractional clearance of urate rather than an over production of urate, occurring as a result of primary hyperuricemia and secondary hyperuricemia. Hyperuricemia is the most important risk factor for developing gout. Gout is the crystal deposition of monosodium urate associated with elevated levels of uric acid. Crystals are deposited in joints, tendons, and surrounding tissues. Some clinical manifestations of gout may include recurrent flares of inflammatory arthritis (gout flare), chronic arthropathy, accumulation of urate crystals in the form of tophaceous deposits, and uric acid nephrolithiasis. Acute attacks of gout are painful and over half of all cases involve the metatarsophalangeal joint of the great toe. Treatment of gout is divided into two phases: acute treatment and chronic prevention. Acute gouty arthritis can be treated with colchicine, NSAIDs, and corticosteroid injections. Urate-lowering agents are uricosuric drugs or xanthine oxidase inhibitors have shown results in reduced frequency of progression of gout to the tophaceous stage. Evidence-based recommendations for the treatment of gout address symptomatic control of acute gout, urate lowering therapy, and prophylaxis of acute attacks. It is recommended to screen patients who are of Chinese, Thai, Korean or other ethnicities who have an increased frequency of the human leukocyte antigen (HLA)-B*5801 gene as giving them allopurinol is associated with an increased risk of severe cutaneous adverse reaction (SCAR), so it is not recommended. Neither allopurinol or febuxostat are recommended in patients concomitantly receiving azathioprine or 6-mercaptopurine, patients with urolithiasis, or those who have a risk of uric acid nephropathy.

Total program savings for the PDL classes will be regularly reviewed.

Program-Specific Information:	Preferred Agents	Non-Preferred Agents
	<ul style="list-style-type: none"> Allopurinol Colchicine Tabs Probenecid Probenecid/Colchicine 	<ul style="list-style-type: none"> Colchicine Caps Colcrys® Febuxostat Gloperba® Mitigare® Uloric® Zyloprim®

Type of Criteria: Increased risk of ADE
 Appropriate Indications

Preferred Drug List
 Clinical Edit

Data Sources: Only Administrative Databases

Databases + Prescriber-Supplied

Setting & Population

- Drug class for review: Antihyperuricemic Agents
- Age range: All appropriate MO HealthNet participants

Approval Criteria

- Failure to achieve desired therapeutic outcomes with trial on 1 or more preferred agents
 - Documented trial period of preferred agents **OR**
 - Documented ADE/ADR to preferred agents **AND**
- For Uloric: adequate therapeutic trial of allopurinol defined as 60 days of therapy in the last 90 days

Denial Criteria

- Lack of adequate trial on required preferred agents
- Therapy will be denied if all approval criteria are not met
- Claim exceeds maximum dosing limitation for the following:

Drug Description	Generic Equivalent	Max Dosing Limitation
ULORIC 40 MG TABLET	FEBUXOSTAT	1 tablet per day
ULORIC 80 MG TABLET	FEBUXOSTAT	1 tablet per day

Required Documentation

Laboratory Results:
MedWatch Form:

Progress Notes:
Other:

Disposition of Edit

Denial: Exception Code "0160" (Preferred Drug List)
Rule Type: PDL

Default Approval Period

1 year

References

- Evidence-Based Medicine Analysis: "Antihyperuricemic Agents", UMKC-DIC; March 2022.
- Evidence-Based Medicine and Fiscal Analysis: "Antihyperuricemic Agents - Therapeutic Class Review", Conduent Business Services, L.L.C., Richmond, VA; June 2021.
- Zylprim [package insert]. East Brunswick, NJ: Casper Pharma; December 2018.
- Colcrys [package insert]. Lexington, MA: Takeda Pharmaceuticals America, Inc.; May 2020.
- Mitigare [package insert]. Memphis, TN: Hikma Americas, Inc.; June 2020.
- Gloperba [package insert]. Alpharetta, GA: Avion Pharmaceuticals, LLC; February 2019.

SmartPA PDL Proposal Form

© 2022 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

Other company trademarks are also acknowledged.

- Uloric [package insert]. Lexington, MA: Takeda Pharmaceuticals America; August 2020.
- Probenecid [package insert]. Parsippany, NJ: Actavis Pharma, Inc.; December 2016.
- Probenecid and colchicine [package insert]. Fairfield, NJ: Ingenus Pharmaceuticals NJ, LLC; 2018.
- Gaffo, A. (2019). Clinical manifestations and diagnosis of gout. In P.L. Romain (Ed.), UpToDate.
- Becker, M., & Perez-Ruiz, F. (2020). Pharmacologic urate-lowering therapy and treatment of tophi in patients with gout. In P.L. Romain (Ed.), UpToDate.
- USPDI, Micromedex; 2022.
- Facts and Comparisons eAnswers (online); 2022 Clinical Drug Information, LLC.

DRAFT