

<b>NEW DRUG REVIEWS FOR 9/17/2010 PA MEETING</b>		
<b>OPEN ACCESS</b>		
<b>FOR INFORMATION ONLY</b>		
<b>COMMON TRADE NAME</b>	<b>GENERIC NAME</b>	<b>INDICATIONS</b>
Amlodipine Besylate 100% Powder	Amlodipine Besylate	Indicated for the treatment of hypertension
Bifera Rx 22-6-1-25 mg Tablet	IRON PS/IRON HEM POLY/FA/B12	Multi-Vitamin
Femcal OB Plus DHA 22-6-1-200 mg Combo Package	Vit D3, Vit E, Vit B1, Vit B2, Vit B3, Vit B12, Folic Acid, Biotin, Iron, DHA	Prenatal Vitamin
Lysteda 650 mcg Tablet	Tranexamic Acid	Indicated for treatment of women suffering from cyclic heavy menstrual bleeding (HMB), also known as menorrhagia
Metvixia 168 mg/G Cream	Methyl Aminolevulinate HCL	Used in combination with red light therapy to treat a skin condition called actinic keratosis of the face and scalp.
Oravig 50 mg MABUC Tablet	Miconazole	Indicated for the treatment of oropharyngeal candidiasis (OPC), more commonly known as thrush, in adults and children age 16 and older
Penderm Cream (GM)	Cream Base No. 34	Emollient cream base
Prenexa 27-1.25-55 mg Capsule	PN/66/iron Fumarate/FA/DSS/DHA	Prenatal Vitamin
Propoven 200 MG/20 ml Ampul	Propofol/PF	For sedation of adults
Rapamune 0.5 mg Tablet	Sirolimus	Indicated for the prophylaxis of organ rejection in patients aged 13years or older receiving renal transplants.
Solu-Cortef 1000 mg/8 ml Vial	Hydrocortisone SOD SUCC/PF	Indicated as follows: Allergic states, Dermatologic diseases, Endocrine disorders, Hematologic disorders, Miscellaneous, Nervous System, Ophthalmic diseases, Renal diseases, Respiratory diseases, Rheumatic disorders.
Solu-Cortef 100 mg/2 ml Vial	Hydrocortisone SOD SUCC/PF	Indicated as follows: Allergic states, Dermatologic diseases, Endocrine disorders, Hematologic disorders, Miscellaneous, Nervous System, Ophthalmic diseases, Renal diseases, Respiratory diseases, Rheumatic disorders.
Solu-Cortef 250 mg/2 ml Vial	Hydrocortisone SOD SUCC/PF	Indicated as follows: Allergic states, Dermatologic diseases, Endocrine disorders, Hematologic disorders, Miscellaneous, Nervous System, Ophthalmic diseases, Renal diseases, Respiratory diseases, Rheumatic disorders.
Solu-Cortef 500 mg/4 ml Vial	Hydrocortisone SOD SUCC/PF	Indicated as follows: Allergic states, Dermatologic diseases, Endocrine disorders, Hematologic disorders, Miscellaneous, Nervous System, Ophthalmic diseases, Renal diseases, Respiratory diseases, Rheumatic disorders.
Torsemide 50 mg/5 ml Vial	Torsemide	Indicated for the treatment of edema associated with congestive heart failure, renal disease, or hepatic disease
Torsemide 20 mg/2 ml	Torsemide	Indicated for the treatment of edema associated with congestive heart failure, renal disease, or hepatic disease
Trelstar 22.5 mg/2 ml Disposable Syringe	Triptorelin Pamoate	Indicated in the palliative treatment of advanced prostate cancer
Trelstar 22.5 mg Vial	Triptorelin Pamoate	Indicated in the palliative treatment of advanced prostate cancer
Tricare Prenatal DHA One 27-1-500 mg Capsule	PNV#20-IRON-FA-DS-FISH-DHA-EPA	Prenatal Vitamin
Zirgan 15% Gel	Ganciclovir	Treatment for acute herpetic keratitis (dendritic ulcers)
Zortress 0.5 mg Tablet	Everolimus	Indicated for the prevention of rejection of kidney transplants in adult patients at low-to-moderate immunologic risk
Zortress 0.75 mg Tablet	Everolimus	Indicated for the prevention of rejection of kidney transplants in adult patients at low-to-moderate immunologic risk
Zortress 0.25 mg Tablet	Everolimus	Indicated for the prevention of rejection of kidney transplants in adult patients at low-to-moderate immunologic risk
Zyclara 3.75% Cream Pack	Imiquimod	Indicated for the topical treatment of clinically typical, visible or palpable actinic keratoses (AK), of the full face or balding scalp in immunocompetent adults. Indicated for the topical treatment of clinically typical, visible or palpable actinic keratoses (AK), of the full face or balding scalp in immunocompetent adults.

<b>PDL EDITS FOR IMPLEMENTATION</b>		
<b>COMMON TRADE NAME</b>	<b>GENERIC NAME</b>	<b>INDICATIONS</b>
Actoplus Met XR 15-1000 mg TBMP 24 Hour	Pioglitazone HCL/Metformin HCL	Indicated for adults with type 2 diabetes who are already treated with ACTOS® (pioglitazone HCl) and metformin or who have inadequate glycemic control on ACTOS or metformin alone <b>NON-PREFERRED</b>
Actoplus Met XR 30-1000 mg TBMP 24 Hour	Pioglitazone HCL/Metformin HCL	Indicated for adults with type 2 diabetes who are already treated with ACTOS® (pioglitazone HCl) and metformin or who have inadequate glycemic control on ACTOS or metformin alone <b>NON-PREFERRED</b>
Differin 0.10% Lotion	Adapalene	Indicated for the treatment of acne <b>NON-PREFERRED</b>
Exalgo ER 8 mg Tablet ER 24 Hour	Hydromorphone HCL	Long Acting Narcotic <b>NON-PREFERRED</b>
Exalgo ER 12 mg Tablet ER 24 Hour	Hydromorphone HCL	Long Acting Narcotic <b>NON-PREFERRED</b>
Exalgo ER 16 mg Tablet ER 24 Hour	Hydromorphone HCL	Long Acting Narcotic <b>NON-PREFERRED</b>
Livalo 1mg Tablet	Pitavastatin Calcium	Indicated as an adjunctive therapy to diet to reduce elevated total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), apolipoprotein B (Apo B), triglycerides (TG), and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia. <b>NON-PREFERRED</b>
Livalo 2mg Tablet	Pitavastatin Calcium	Indicated as an adjunctive therapy to diet to reduce elevated total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), apolipoprotein B (Apo B), triglycerides (TG), and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia. <b>NON-PREFERRED</b>
Livalo 4mg Tablet	Pitavastatin Calcium	Indicated as an adjunctive therapy to diet to reduce elevated total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), apolipoprotein B (Apo B), triglycerides (TG), and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia. <b>NON-PREFERRED</b>
Prolia 60 mg/ml Disposabe Syringe	Denosumab	Indicated to treat osteoporosis in patients at high risk of fracture <b>NON-PREFERRED</b>
Rybix ODT 50 mg Tablet Rapid Dissolve	Tramadol HCL	Indicated for the management of moderate to moderately severe pain in adults <b>NON-PREFERRED</b>
Zymaxid 0.5% Drops	gatifloxacin ophthalmic 0.5% solution	A topical fluoroquinolone anti-infective indicated for the treatment of bacterial conjunctivitis caused by susceptible strains <b>NON-PREFERRED</b>
<b>CLINICAL EDITS FOR IMPLEMENTATION</b>		
<b>COMMON TRADE NAME</b>	<b>GENERIC NAME</b>	<b>INDICATIONS</b>
Cambia 50 mg Powder Pack	Diclofenac Potassium	Used to treat migraine attacks in adults
Vimpat 10 mg/ml Solution	Lacosamide	15 Day Supply Edit
Vimovo 375 mg/20 mg TBMP 12 Hour	Naproxen/Someprazole Mag	Indications are for the signs and symptoms of osteoarthritis, rheumatoid arthritis and ankylosing spondylitis in patients who are at risk for developing NSAID-associated ulcers.
Vimovo 500 mg/20 mg TBMP 12 Hour	Naproxen/Someprazole Mag	Indications are for the signs and symptoms of osteoarthritis, rheumatoid arthritis and ankylosing spondylitis in patients who are at risk for developing NSAID-associated ulcers.
<b>REQUEST FOR CONTINUED PRIOR AUTHORIZATION</b>		
<b>COMMON TRADE NAME</b>	<b>GENERIC NAME</b>	<b>INDICATIONS</b>
Cenhist 15 mg-6 mg Chewable Tablet	Phenylephrine/Brompheniramine	Cough and Cold
Novaferum 100-1-15/5 ml Soution Reconstituted	IRON PS CMPLX/FA/VIT B12/VIT C	Multi-vitamin
Orbivan 50-300-40 mg Capsule	BUTALB/ACETAMINOPHEN/CAFFEINE	Indicated for the relief of the symptom complex of tension (or muscle contraction) headache
Pennsaid 1.50% Solution	Diclofenac Sodium	Used for the treatment of the signs and symptoms of osteoarthritis of the knee.
Sodium Sulfacetamide Sulfur 9%-4% Cleanser	Sulfacetamide Sodium/Sulfur	Used in the treatment of acne, rosacea, and seborrheic dermatitis (a red, flaking skin rash)
Xifaxan 550 mg Tablet	Rifaximin	Antibiotic that fights bacterial infection only in the intestines. It is used to treat travelers' diarrhea caused by E. coli. Most people get this infection by eating food or drinking fluids that have been contaminated with the bacteria.