



# SmartPA Criteria Proposal

Drug/Drug Class:	Opioids – Short-Acting Single Agents Clinical Edit	
First Implementation Date:	February 8, 2012	
Proposed Date:	March 18, 2021	
Prepared for:	MO HealthNet	
Prepared by:	MO HealthNet/Conduent	
Criteria Status:	□Existing Criteria ☑Revision of Existing Criteria □New Criteria	

#### **Executive Summary**

Purpose: Ensure safe, appropriate utilization and control of short-acting single agent opioids

Why Issue Selected:

Short-acting single agent opioids are indicated for short-term relief of moderate to severe pain on an "as needed" basis. These medications are often used in conjunction with a long acting opioids to help relieve breakthrough pain. Although highly effective for pain control, these substances carry with them the potential for harm from adverse drug events and/or overdose. Opioid drugs also have substantial misuse liability and are often implicated among persons who have developed a substance misuse disorder. Concomitant use of benzodiazepines significantly increases the risk of harm from opioids.

In March 2016, the Centers for Disease Control and Prevention (CDC) released their Guideline for Prescribing Opioids for Chronic Pain. Although these guidelines focus on managing chronic pain, they emphasize the need for prescribers to be more judicious in the initiation, continuation, selection, and monitoring of opioids and other medications for acute pain. As part of the efforts to protect participants from the possible adverse effects of opioid medications and subsequent diversion or misuse of opioid medications, MO HealthNet Division will continue to clinically edit the use of these controlled substances. Limiting or reducing "morphine equivalent doses" helps avoid harmful effects of opioids and promotes patient safety. Opioid daily doses above 50 MME (morphine milligram equivalents) increase the risk of overdose by at least double. Other preventative activities include limiting prescription amounts and frequency of dispensing.

In a Safety Announcement dated April 20, 2017, the Food and Drug Administration (FDA) advised that it is restricting the use of codeine and tramadol medicines in children. Codeine is approved to treat pain and cough, and tramadol is approved to treat pain. These medicines carry serious risks, including slowed or difficult breathing and death, which appear to impose a greater risk in children younger than 12 years, and hence codeine and tramadol containing agents should not be used in these children. These medicines should also be limited in some older children. Single-ingredient codeine and all tramadol-containing products are FDA-approved only for use in adults. The FDA is also recommending against the use of codeine and tramadol medicines in breastfeeding mothers due to possible harm to their infants.

Type of Criteria:	<ul><li>☑ Increased risk of ADE</li><li>☑ Appropriate Indications</li></ul>	<ul><li>□ Preferred Drug List</li><li>☑ Clinical Edit</li></ul>
Data Sources:	☐ Only Administrative Databases	□ Databases + Prescriber-Supplied

#### **Setting & Population**

- Drug class for review: Short-acting single agent opioids
- Age range: All appropriate MO HealthNet participants

#### **Approval Criteria**

- Documented diagnosis of cancer in the past 6 months OR
- Documented diagnosis of sickle cell disease in the past 6 months OR
- Participant currently enrolled in Hospice care or receiving palliative care in the past year OR
- Claim does not exceed approved dosage limitations:
  - Participant aged < 18 years: see Appendix B OR</li>
  - Participant aged ≥ 18 years: see Appendix A OR
  - Participant demonstrates compliance without dose escalation to prescribed therapy over the current MME threshold
- Claim is for methadone and participant has no prior history of methadone therapy in the past 90 days: Clinical Consultant Review required OR
- If no prior history of opioid therapy in the past 90 days:
  - Claim is for ≤ 7 days supply AND
  - Claim is for ≤ 50 MME per day (see Appendix C) OR
- If claim plus history equals > 60 days of short acting opioid therapy in the past 90 days: documented diagnosis of chronic non-malignant pain (CNMP) in the past 6 months required OR
- Approval based upon clinical consultant review and/or receipt of signed Opioid Attestation Form from prescriber

#### **Denial Criteria**

- Therapy will be denied if all approval criteria are not met
- Documented opioid dependence therapy in the past 45 days
- Participant aged < 12 years and claim is for a codeine agent</li>
- Prior history of any short acting opioid in the past 3 days
- Denial criteria contained within the High Risk Therapies Clinical Edit: Claim is for an opioid (excluding buprenorphine tablets and buprenorphine/naloxone combinations) and:
  - Participant has history of > 3 days of select oral benzodiazepine therapy (alprazolam, chlordiazepoxide, clonazepam, clorazepate, diazepam, lorazepam, and oxazepam) in the past 60 days AND
  - Participant lacks history of at least 1 claim for an opioid emergency reversal agent in the past 2 years

Required Documenta	tion			
Laboratory Results: MedWatch Form:		Progress Notes: Other:	X	

SmartPA Clinical Proposal Form

© 2021 Conduent Business Services, LLC. All rights reserved. Conduent<sup>TM</sup> and Conduent Design<sup>TM</sup> are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

## **Disposition of Edit**

Denial: Exception code "0682" (Clinical Edit)

Rule Type: CE

## **Default Approval Period**

7 days

### Appendix A Adult Dose Chart

Drug Description	Max Units per Day
BUTORPHANOL TARTRATE 10 MG/ML NASAL SPRAY	0.5 ML
CODEINE SULFATE 15 MG ORAL TABLET	24 TABLETS
CODEINE SULFATE 30 MG ORAL TABLET	12 TABLETS
CODEINE SULFATE 60 MG ORAL TABLET	6 TABLETS
DSUVIA 30 MCG SL TAB	12 TABLETS
HYDROMORPHONE 1 MG/ML AMPULE	22 ML
HYDROMORPHONE 2 MG/ML AMPULE	11 ML
HYDROMORPHONE 4 MG/ML AMPULE	5.5 ML
HYDROMORPHONE HYDOCHLORIDE 1 MG/ML SOL	22 ML
HYDROMORPHONE HYDROCHLORIDE 2 MG TABLET	11 TABLETS
HYDROMORPHONE HYDROCHLORIDE 3 MG RECTAL SUPP	4 SUPPOSITORIES
HYDROMORPHONE HYDROCHLORIDE 4 MG TABLET	5.5 TABLETS
HYDROMORPHONE HYDROCHLORIDE 8 MG TABLET	2.5 TABLETS
LEVORPHANOL 2 MG TABLET	12 TABLETS
LEVORPHANOL 3 MG TAB	8 TABLETS
MEPERIDINE HYDROCHLORIDE 100 MG ORAL TAB	6 TABLETS
MEPERIDINE HYDROCHLORIDE 50 MG ORAL TABLET	12 TABLETS
MEPERIDINE HYDROCHLORIDE 50 MG/5 ML ORAL SOLUTION	60 ML
METHADONE HYDROCHLORIDE 10 MG TABLET	-
METHADONE HYDROCHLORIDE 10 MG/5 ML SOL	-
METHADONE HYDROCHLORIDE 10 MG/ML CONC SOL	-
METHADONE HYDROCHLORIDE 40 MG ODT	-
METHADONE HYDROCHLORIDE 5 MG TABLET	-
METHADONE HYDROCHLORIDE 5 MG/5 ML SOL	-
MORPHINE SULF 10MG/0.5ML SOLN	4.5 ML
MORPHINE SULFATE 10 MG RECTAL SUPPOSITORY	6 SUPPOSITORIES
MORPHINE SULFATE 10 MG/5 ML SOLUTION	45 ML
MORPHINE SULFATE 100 MG/5 ML SOLUTION	4.5 ML
MORPHINE SULFATE 15 MG IMMEDIATE-RELEASE TAB	6 TABLETS
MORPHINE SULFATE 20 MG RECTAL SUPPOSITORY	4 SUPPOSITORIES
MORPHINE SULFATE 20 MG/5 ML SOLUTION	22.5 ML
MORPHINE SULFATE 30 MG IMMEDIATE-RELEASE TAB	3 TABLETS
MORPHINE SULFATE 30 MG RECTAL SUPPOSITORY	3 SUPPOSITORIES
MORPHINE SULFATE 5 MG RECTAL SUPPOSITORY	6 SUPPOSITORIES
NALBUPHINE 10 MG/0.5 ML SYR	8 ML
NALBUPHINE 10 MG/ML AMPUL	16 ML
NALBUPHINE 100 MG/10 ML VIAL	16 ML
NALBUPHINE 20 MG/ML AMPUL	8 ML
NALBUPHINE 200 MG/10 ML VIAL	8 ML

SmartPA Clinical Proposal Form

© 2021 Conduent Business Services, LLC. All rights reserved. Conduent<sup>TM</sup> and Conduent Design<sup>TM</sup> are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

OXYCODONE 10 MG/0.5 ML ORAL SYR	3 ML
OXYCODONE HYDROCHLORIDE 10 MG ORAL TABLET	6 TABLETS
OXYCODONE HYDROCHLORIDE 15 MG ORAL TABLET	4 TABLETS
OXYCODONE HYDROCHLORIDE 20 MG ORAL TABLET	3 TABLETS
OXYCODONE HYDROCHLORIDE 20 MG/ML ORAL SOL	3 ML
OXYCODONE HYDROCHLORIDE 30 MG ORAL TABLET	2 TABLETS
OXYCODONE HYDROCHLORIDE 5 MG ORAL CAPSULE	12 CAPSULES
OXYCODONE HYDROCHLORIDE 5 MG ORAL TABLET	12 TABLETS
OXYCODONE HYDROCHLORIDE 5 MG/5 ML ORAL SOL	60 ML
OXYCODONE HYDROCHLORIDE 7.5 MG ORAL TABLET	8 TABLETS
OXYMORPHONE HYDROCHLORIDE 10 MG ORAL TAB	3 TABLETS
OXYMORPHONE HYDROCHLORIDE 5 MG ORAL TAB	6 TABLETS
ROXYBOND 15 MG ORAL TAB	4 TABLETS
ROXYBOND 30 MG ORAL TAB	2 TABLETS

## Appendix B Pediatric Dose Chart

Drug Description	Max Units per Day
CODEINE SULFATE 15 MG ORAL TABLET	24 TABLETS
CODEINE SULFATE 30 MG ORAL TABLET	12 TABLETS
CODEINE SULFATE 60 MG ORAL TABLET	6 TABLETS
HYDROMORPHONE 1 MG/ML AMPULE	12 ML
HYDROMORPHONE 2 MG/ML AMPULE	6 ML
HYDROMORPHONE 4 MG/ML AMPULE	3 ML
HYDROMORPHONE HYDOCHLORIDE 1 MG/ML SOLUTION	12 ML
HYDROMORPHONE HYDROCHLORIDE 2 MG TABLET	6 TABLETS
MORPHINE SULFATE 10 MG RECTAL SUPPOSITORY	6 SUPPOSITORIES
MORPHINE SULF 10MG/0.5ML SOLN	3 ML
MORPHINE SULFATE 10 MG/5 ML SOLUTION	30 ML
MORPHINE SULFATE 100 MG/5 ML SOLUTION	4 ML
MORPHINE SULFATE 15 MG IMMEDIATE-RELEASE TAB	2 TABLETS
MORPHINE SULFATE 20 MG/5 ML SOLUTION	15 ML
MORPHINE SULFATE 5 MG RECTAL SUPPOSITORY	6 SUPPOSITORIES
OXYCODONE 10 MG/0.5 ML ORAL SYR	1.5 ML
OXYCODONE HYDROCHLORIDE 10 MG ORAL TABLET	3 TABLETS
OXYCODONE HYDROCHLORIDE 15 MG ORAL TABLET	2 TABLETS
OXYCODONE HYDROCHLORIDE 20 MG/ML ORAL SOL	1.5 ML
OXYCODONE HYDROCHLORIDE 5 MG ORAL CAPSULE	6 CAPSULES
OXYCODONE HYDROCHLORIDE 5 MG ORAL TABLET	6 TABLETS
OXYCODONE HYDROCHLORIDE 5 MG/5 ML ORAL SOL	30 ML
OXYCODONE HYDROCHLORIDE 7.5 MG ORAL TABLET	4 TABLETS
ROXYBOND 15 MG ORAL TAB	2 TABLETS
ROXYBOND 30 MG ORAL TAB	1 TABLET

#### Opioid Naïve Dose Chart Appendix C

Drug Description	Max Units per Day	
BUTORPHANOL TARTRATE 10 MG/ML NASAL SPRAY	0.5 ML	
CODEINE SULFATE 15 MG ORAL TABLET	22 TABLETS	
CODEINE SULFATE 30 MG ORAL TABLET	11 TABLETS	
CODEINE SULFATE 60 MG ORAL TABLET	5.5 TABLETS	
DSUVIA 30 MCG SL TAB	12 TABLETS	

SmartPA Clinical Proposal Form

© 2021 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

HYDROMORPHONE 1 MG/ML AMPULE	12 ML
HYDROMORPHONE 2 MG/ML AMPULE	6 ML
HYDROMORPHONE 4 MG/ML AMPULE	3 ML
HYDROMORPHONE HYDOCHLORIDE 1 MG/ML SOL	12 ML
HYDROMORPHONE HYDROCHLORIDE 2 MG TABLET	6 TABLETS
HYDROMORPHONE HYDROCHLORIDE 3 MG RECTAL SUPP	4 SUPPOSITORIES
HYDROMORPHONE HYDROCHLORIDE 3 MIG RECTAL SUPP	3 TABLETS
HYDROMORPHONE HYDROCHLORIDE 8 MG TABLET	1.5 TABLETS
LEVORPHANOL 2 MG TABLET	12 TABLETS
LEVORPHANOL 3 MG TAB	8 TABLETS
MEPERIDINE HYDROCHLORIDE 100 MG ORAL TAB	5 TABLETS
MEPERIDINE HYDROCHLORIDE 50 MG ORAL TAB	10 TABLETS
MEPERIDINE HYDROCHLORIDE 50 MG/5 ML ORAL SOLUTION	50 ML
METHADONE HYDROCHLORIDE 10 MG TABLET	1 TABLETS
METHADONE HYDROCHLORIDE 10 MG/5 ML SOL	5 ML
METHADONE HYDROCHLORIDE 10 MG/ML CONC SOL	1 ML
METHADONE HYDROCHLORIDE 40 MG ODT	0.25 TABLETS
METHADONE HYDROCHLORIDE 5 MG TABLET	2 TABLETS
METHADONE HYDROCHLORIDE 5 MG/5 ML SOL	10 ML
MORPHINE SULF 10MG/0.5ML SOLN	2.5 ML
MORPHINE SULFATE 10 MG RECTAL SUPPOSITORY	5 SUPPOSITORIES
MORPHINE SULFATE 10 MG/5 ML SOLUTION	25 ML
MORPHINE SULFATE 100 MG/5 ML SOLUTION	2.5 ML
MORPHINE SULFATE 15 MG IMMEDIATE-RELEASE TAB	3 TABLETS
MORPHINE SULFATE 20 MG RECTAL SUPPOSITORY	2 SUPPOSITORIES
MORPHINE SULFATE 20 MG/5 ML SOLUTION	12.5 ML
MORPHINE SULFATE 30 MG IMMEDIATE-RELEASE TAB	1.5 TABLETS
MORPHINE SULFATE 30 MG RECTAL SUPPOSITORY	1.5 SUPPOSITORIES
MORPHINE SULFATE 5 MG RECTAL SUPPOSITORY	6 SUPPOSITORIES
NALBUPHINE 10 MG/0.5 ML SYR	8 ML
NALBUPHINE 10 MG/ML AMPUL	16 ML
NALBUPHINE 100 MG/10 ML VIAL	16 ML
NALBUPHINE 20 MG/ML AMPUL	8 ML
NALBUPHINE 200 MG/10 ML VIAL	8 ML
OXYCODONE 10 MG/0.5 ML ORAL SYR	1.5 ML
OXYCODONE HYDROCHLORIDE 10 MG ORAL TAB	3 TABLETS
OXYCODONE HYDROCHLORIDE 15 MG ORAL TABLET	2 TABLETS
OXYCODONE HYDROCHLORIDE 20 MG ORAL TABLET	1.5 TABLETS
OXYCODONE HYDROCHLORIDE 20 MG/ML ORAL SOL	1.5 ML
OXYCODONE HYDROCHLORIDE 30 MG ORAL TABLET	1 TABLETS
OXYCODONE HYDROCHLORIDE 5 MG ORAL CAP	6 CAPSULES
OXYCODONE HYDROCHLORIDE 5 MG ORAL TABLET	6 TABLETS
OXYCODONE HYDROCHLORIDE 5 MG/5 ML ORAL SOL	30 ML
OXYCODONE HYDROCHLORIDE 7.5 MG ORAL TABLET	4 TABLETS
OXYMORPHONE HYDROCHLORIDE 10 MG ORAL TAB	1.5 TABLETS
OXYMORPHONE HYDROCHLORIDE 5 MG ORAL TAB	3 TABLETS
ROXYBOND 15 MG ORAL TAB	2 TABLETS
ROXYBOND 30 MG ORAL TAB	1 TABLET
TOTT DOING ONAL TAD	IIADELI

## References

SmartPA Clinical Proposal Form

© 2021 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

- Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain United States, 2016. MMWR Recomm Rep 2016;65(No. RR-1):1–49. DOI: <a href="http://dx.doi.org/10.15585/mmwr.rr6501e1">http://dx.doi.org/10.15585/mmwr.rr6501e1</a>
- U.S. Food and Drug Administration. Drug Safety Communications: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. <a href="https://www.fda.gov/media/104268/download">https://www.fda.gov/media/104268/download</a>. Accessed November 9, 2020.
- Oregon Health Authority Opioid Conversion Calculator.
   <a href="https://www.oregonpainguidance.org/opioidmedcalculator/">https://www.oregonpainguidance.org/opioidmedcalculator/</a>. Accessed November 9, 2020.
- Facts and Comparisons. Opioid Analgesics. Accessed November 9, 2020.