

SmartPA Criteria Proposal

Drug/Drug Class:	Benign Prostatic Hyperplasia Agents PDL Edit
First Implementation Date:	November 14, 2007
Proposed Date:	June 18, 2020
Prepared For:	MO HealthNet
Prepared By:	MO HealthNet/Conduent
Criteria Status:	<input checked="" type="checkbox"/> Existing Criteria <input type="checkbox"/> Revision of Existing Criteria <input type="checkbox"/> New Criteria

Executive Summary

Purpose: The MO HealthNet Pharmacy Program will implement a state-specific preferred drug list.

Why Issue Selected: In adult men, the prostate acts mainly to add fluids to the semen that nourish sperm and protect them against the natural acids in the vagina. Benign prostatic hyperplasia (BPH) is one of the most common diseases in aging men – for most men over age 50. The symptoms are brought on by hyperplastic changes in the prostate, leading to prostatic enlargement. The resulting urinary obstruction increases outflow resistance and results in a detrusor muscle response. These participants often present with bothersome lower urinary tract symptoms, including frequency, nocturia, urgency, and urge incontinence resulting from irritation; and/or symptoms secondary to obstruction, such as difficulty initiating urination, or passing urine, weak stream, involuntary post-void dripping of urine, or a sensation of incomplete bladder emptying. Unless participants have developed bladder outlet obstruction, BPH only requires therapy if symptoms have a significant impact on a participant's quality of life. The agents that are most commonly used to treat the lower urinary tract symptoms associated with BPH are alpha-1 adrenergic antagonists, 5-alpha-reductase inhibitors, anticholinergic agents, and phosphodiesterase-5 inhibitors. In mild to moderate symptomatic participants, an alpha-1 adrenergic antagonist as monotherapy is recommended. In severe symptomatic participants, a combination of an alpha-1 adrenergic antagonist and a 5-alpha-reductase inhibitor is recommended. Alpha-1 adrenergic antagonists work in the lower urinary tract by blocking adrenoreceptors in the prostate to cause smooth muscles to relax and thus improve urine flow rate and reduce BPH symptoms. Selective 5-alpha reductase inhibitors decrease the serum concentration of 5a-dihydrotestosterone (DHT) which stimulates the growth of glandular and stromal cells, increasing prostatic tissue mass.

Total program savings for the PDL classes will be regularly reviewed.

Program-Specific Information:	Non-Preferred Agents
<ul style="list-style-type: none"> • Alfuzosin • Doxazosin • Dutasteride • Finasteride 5mg • Prazosin • Tamsulosin • Terazosin 	<ul style="list-style-type: none"> • Avodart® • Cardura® • Cardura® XL • Cialis® 5mg • Dutasteride/Tamsulosin • Flomax® • Jayln® • Minipress® • Proscar® • Rapaflo® • Silodosin • Tadalafil 5mg • Uroxatral®

- Type of Criteria:** Increased risk of ADE Preferred Drug List
 Appropriate Indications Clinical Edit
- Data Sources:** Only Administrative Databases Databases + Prescriber-Supplied

Setting & Population

- Drug class for review: Benign Prostatic Hyperplasia Agents
- Age range: All appropriate MO HealthNet participants

Approval Criteria

- Failure to achieve desired therapeutic outcomes with trial on 3 or more preferred agents
 - Documented trial period of preferred agents **OR**
 - Documented ADE/ADR to preferred agents **AND**
- For Cialis: documented diagnosis of benign prostatic hyperplasia in the past 2 years

Denial Criteria

- Lack of adequate trial on required preferred agents
- Therapy will be denied if no approval criteria are met
- For Cialis:
 - Documented diagnosis of erectile dysfunction in the past 2 years **OR**
 - Claim history documents use of nitrates or ritonavir therapy in the past 30 days **OR**
 - Documented contraindication to tadalafil:
 - History of MI in the past 90 days
 - History of unstable angina
 - History of NYHA Class II or greater heart failure
 - History of stroke in the past 6 months
 - History of uncontrolled arrhythmias

Required Documentation

Laboratory Results: Progress Notes:
 MedWatch Form: Other:

SmartPA PDL Proposal Form
 © 2020 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.
 Other company trademarks are also acknowledged.

Disposition of Edit

Denial: Exception Code "0160" (Preferred Drug List)
Rule Type: PDL

Default Approval Period

1 year

References

1. Lippincott, Williams, Wilkins. PDR Electronic Library, Montvale NJ; 2020.
2. USPDI, Micromedex; 2020.
3. Facts and Comparisons eAnswers (online); 2020 Clinical Drug Information, LLC.
4. Cunningham, G., & Kadmon, D., (2019). Medical treatment of benign prostatic hyperplasia. In J. Givens (Ed.), *UptoDate*.
5. Evidence-Based Medicine and Fiscal Analysis: "Benign Prostatic Hyperplasia (BPH) Agents – Therapeutic Class Review", Conduent Business Services, L.L.C., Richmond, VA; April 2020.
6. Evidence-Based Medicine Analysis: "BPH Inhibitors", UMKC-DIC; February 2020.
7. Terazosin [package insert]. Pulaski, TN: AvKARE; 2020.
8. Uroxatral [package insert]. St. Michael, Barbados: Concordia Pharmaceuticals, Inc; 2019.
9. Proscar [package insert]. Whitehouse Station, NJ: Merck & Co; 2013.
10. Jalyn [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2017.
11. Avodart [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2020.
12. Cialis [package insert]. Indianapolis, IN: Eli Lilly and Company; 2018.
13. Cardura [package insert]. New York, NY: Pfizer; 2019.
14. Cardura XL [package insert]. New York, NY: Pfizer; 2017.
15. Flomax [package insert]. Bridgewater, NJ: Sanofi-Aventis US LLC; 2019.
16. Rapaflo [package insert]. Irvine, CA; Allergan USA, Inc; 2017.

SmartPA PDL Proposal Form

© 2020 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

Other company trademarks are also acknowledged.