



# SmartPA Criteria Proposal

Drug/Drug Class:	Narcolepsy Inhibitors Clinical Edit	
First Implementation Date:	August 27, 2013	
Proposed Date:	September 17, 2020	
Prepared for:	MO HealthNet	
Prepared by:	MO HealthNet/Conduent	
Criteria Status:	□Existing Criteria ☑Revision of Existing Criteria □New Criteria	

#### **Executive Summary**

Purpose: Ensure appropriate utilization and control of narcolepsy inhibitors

### Why Issue Selected:

Narcolepsy is a chronic sleep disorder characterized by excessive daytime sleepiness and a neurological inability to regulate sleep-wake cycles. It affects an estimated 1 in 2,000 people in the United States. Excessive Daytime Sleepiness (EDS) is the primary symptom of narcolepsy, characterized by the inability to stay awake and alert during the day and resulting in unplanned lapses into sleep or drowsiness. The sleepiness may be so severe that patients with narcolepsy can rapidly doze off with little warning. Other primary symptoms include cataplexy, sleep related hallucinations, sleep paralysis and sleep disruption. While all patients with narcolepsy experience excessive daytime sleepiness, they may not experience all primary symptoms. Obstructive Sleep Apnea (OSA) is the most common sleep-related breathing disorder and is also associated with excessive daytime sleepiness. The estimated prevalence of OSA in the United States is approximately 15% in males and 5% in females. Although the underlying cause is different, EDS in OSA is treated similarly to EDS due to narcolepsy.

Provigil® (modafinil) and Nuvigil® (armodafinil) are indicated to improve wakefulness in adult patients with excessive sleepiness associated with OSA, narcolepsy, or shift work disorder; these first-line therapies have been widely and safely used since 1998. Xyrem® (sodium oxybate) was FDA approved in 2002 and is indicated for the treatment of cataplexy or excessive daytime sleepiness in patients ≥ 7 years old with narcolepsy. Xyrem is also known as the "date rape" drug because it can cause decreased consciousness; it has a REMS program due to risks of CNS depression and abuse. Recently FDA approved in March 2019, Sunosi® (solriamfetol) is indicated to improve wakefulness in adult patients with excessive daytime sleepiness associated with narcolepsy or OSA. Also, FDA approved in August 2019, Wakix® (pitolisant) is indicated for treatment of excessive daytime sleepiness in adult patients with narcolepsy; Wakix is the first H3 receptor antagonist and the first non-controlled substance approved to treat narcolepsy.

On July 22, 2020, the FDA approved Xywav<sup>™</sup> (calcium, magnesium, potassium, and sodium oxybates), a low-sodium alternative to Xyrem. Xywav is not expected to be launched until late in 2020 and is not currently included in this edit.

## Program-Specific Information:

Date Range FFS 7-1-2019 to 06-30-2020					
Drug	Claims	Spend	Cost per unit (tablet/ml)	Avg spend per claim	
NUVIGIL 50 MG TAB	37	\$1,047.17	\$0.31 NADAC	\$28.30	
NUVIGIL 150 MG TAB	236	\$11,499.39	\$0.92 NADAC	\$48.72	
NUVIGIL 200MG TAB	54	\$2,058.05	\$0.90 NADAC	\$38.11	
NUVIGIL 250 MG TAB	364	\$18,828.26	\$0.92 NADAC	\$51.72	
PROVIGIL 100 MG TAB	340	\$12,073.13	\$0.48 NADAC	\$35.50	
PROVIGIL 200 MG TAB	921	\$39,756.87	\$0.63 NADAC	\$43.16	
SUNOSI 75MG TAB	9	\$5,428.77	\$21.12 NADAC	\$603.19	
SUNOSI 150MG TAB	14	\$8,729.50	\$21.16 NADAC	\$623.53	
WAKIX 4.45 MG TAB	3	\$4,008.15	\$94.75 WAC	\$1,336.05	
WAKIX 17.8 MG TAB	15	\$107,210.75	\$189.50 WAC	\$7,147.38	
XYREM 500 MG/ML SOLN	155	\$2,057,885.19	\$28.27 MAC	\$13,276.67	

Type of Criteria.	M increased risk of ADE	Preferred Drug List	

Data Sources: ☐ Only Administrative Databases ☐ Databases + Prescriber-Supplied

#### **Setting & Population**

- Drug class for review: Narcolepsy inhibitors
- Age range: All appropriate MO HealthNet participants aged 7 years and older

#### **Approval Criteria**

- Documented compliance to current therapy without concurrent stimulant use (90 days in the past 120 days) OR
- For modafinil and armodafinil:
  - o Participant age ≥ 16 years for modafanil AND
  - Participant age ≥ 17 years for armodafinil AND
  - Documented diagnosis of one of the following:
    - Obstructive sleep apnea with history of CPAP
    - Shift work disorder
    - Narcolepsy
    - Hypersomnia
    - Fatigue related to multiple sclerosis
- For Sunosi (solriamfetol):
  - o Participant age ≥ 18 years AND
  - Documented trial of modafinil or armodafinil in the past year AND
  - Documented trial of a stimulant in the past year AND
  - Documented diagnosis of obstructive sleep apnea with history of CPAP OR
  - Documented diagnosis of narcolepsy
- For Wakix (pitolisant):
  - Participant age ≥ 18 years AND
  - Documented diagnosis of narcolepsy AND
  - Documented trial of modafinil or armodafinil in the past year AND
  - Documented trial of a stimulant in the past year AND
  - Documented trial of solriamfetol in the past year
- For sodium oxybate:
  - Documented diagnosis of narcolepsy with cataplexy and participant age ≥ 7 years OR

SmartPA Clinical Proposal Form

© 2020 Conduent Business Services, LLC. All rights reserved. Conduent $^{TM}$  and Conduent Design $^{TM}$  are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

- Documented diagnosis of narcolepsy with excessive daytime sleepiness
  - Participant age ≥ 7 years and < 18 years: documented trial of a stimulant in the past year</li>
  - Participant age ≥ 18 years:
    - Documented trial of modafinil or armodafinil in the past year AND
    - Documented trial of a stimulant in the past year AND
    - Documented trial of solriamfetol in the past year AND
    - Documented trial of pitolisant in the past year

#### **Denial Criteria**

- Therapy will be denied if all approval criteria are not met
- Claim exceeds daily dosing limitations (see Appendix A)
- For sodium oxybate:
  - o Documented history of substance abuse OR
  - Documented history of renal impairment OR
  - Documented history of heart failure OR
  - Documented history of uncontrolled hypertension in the past year OR
  - Documented history of suicide attempt
- For Wakix (pitolisant): Documented diagnosis of severe hepatic impairment

<b>Required Documentatio</b>	n		
Laboratory Results: MedWatch Form:	Progress Notes: Other:	X	
Disposition of Edit			
Denial: Exception code "068 Rule Type: CE	32" (Clinical Edit)		
Default Approval Period			

### Appendix A

1 year

Drug Description	Generic Equivalent	Max Units per Day
Provigil 100 mg tablet	modafinil	2 tablets
Provigil 200 mg tablet	modafinil	2 tablets
Nuvigil 50 mg tablet	armodafinil	1 tablet
Nuvigil 150 mg tablet	armodafinil	1 tablet
Nuvigil 200 mg tablet	armodafinil	1 tablet
Nuvigil 250 mg tablet	modafinil	1 tablet
Xyrem 500 mg/ml solution	sodium oxybate	18 milliliters
Sunosi 75mg tablet	solriamfetol	1 tablet
Sunosi 150mg tablet	solriamfetol	1 tablet
Wakix 4.45mg tablet	pitolisant	2 tablets
Wakix 17.8mg tablet	pitolisant	2 tablets

SmartPA Clinical Proposal Form

© 2020 Conduent Business Services, LLC. All rights reserved. Conduent $^{TM}$  and Conduent Design $^{TM}$  are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

#### References

- PROVIGIL® (modafinil) [package insert]. North Wales, PA: Teva Pharmaceuticals USA, Inc.; July 2019.
- NUVIGIL® (armodafinil) [package insert]. North Wales, PA: Teva Pharmaceuticals USA, Inc.; July 2019.
- XYREM® (sodium oxybate) [package insert]. Palo Alto, CA: Jazz Pharmaceuticals, Inc.; July 2020.
- SUNOSI® (solriamfetol) [package insert]. Palo Alto, CA: Jazz Pharmaceuticals, Inc.; June 2019.
- WAKIX® (pitolisant) [package insert]. Plymouth Meeting, PA: Harmony Biosciences, LLC; November 2019.
- IPD Analytics. CNS: Sleep Disorders. Accessed August 4, 2020.
- IPD Analytics. New Drug Approval: Sunosi (solriamfetol). May 2019.
- IPD Analytics. New Drug Approval: Wakix (pitolisant). September 2019.
- IPD Analytics. Jazz's Xywav Approved for Narcolepsy as Low-Sodium Alternative to Xyrem. July 27, 2020.

