



## Missouri Pharmacy Program – Preferred Drug List



### Tramadol-Like Agents

Effective 06/21/2012

Revised 04/05/2018

#### Preferred Agents

- Tramadol Tablets
- Tramadol-Acetaminophen (APAP) Tablets
- **Tramadol ER**

#### Non-Preferred Agents

- Conzip® ER Capsules
- Nucynta® Tablets
- Nucynta ER® Tablets
- Ryzolt ER®/ODT Tablets
- Synapryn FusePaq® Susp
- Ultracet® Tablets
- Ultram® Tablets
- Ultram ER® Tablets

### Approval Criteria

- Failure to achieve desired therapeutic outcomes with trial on 2 preferred agents
  - Documented trial period for preferred agents
  - Documented ADE/ADR to preferred agents
- Documented compliance on current therapy regimen

### Denial Criteria

#### Ultram:

- **Under 12 years of age**
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\leq$  75 years of age – 400 mg
  - Patients  $>$  75 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Ultracet:**

- **Under 12 years of age**
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  16 years of age – 300 mg tramadol and 2600mg acetaminophen
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Ultram ER/Ryzolt:**

- **Under 12 years of age**
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  16 years of age – 300 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days
- History of opioid abuse or dependence – therapy will be subject to Clinical Consultant review
- Claim for Ultram ER/Ryzolt therapy without previous stabilization on immediate-release tramadol product.

**Nucynta:**

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  18 years of age – 700 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days

**Nucynta ER:**

- 17 years of age or less
- Daily dose exceeds maximum recommended daily dose:
  - Patients  $\geq$  18 years of age – 500 mg
- Presence of seizure history
- Concurrent use of monoamine-oxidase inhibitor or discontinuation of one within 14 days