



SmartPA Criteria Proposal

Drug/Drug Class:	Anticonvulsants, Rescue Agents PDL Edit
First Implementation Date:	January 21, 2021
Revised Date:	January 12, 2023
Prepared For:	MO HealthNet
Prepared By:	MO HealthNet/Conduent
Criteria Status:	<input type="checkbox"/> Existing Criteria <input checked="" type="checkbox"/> Revision of Existing Criteria <input type="checkbox"/> New Criteria

Executive Summary

Purpose: The MO HealthNet Pharmacy Program will implement a state-specific preferred drug list.

Why Issue Selected: There are multiple FDA approved agents for the treatment of acute repetitive seizures or clusters. Diazepam rectal (Diastat®) is indicated for the management of select, refractory participants 2 years of age or older with epilepsy on stable regimens of antiepileptic drugs, who require intermittent use of diazepam to control episodes of increased seizure activity. Diazepam intranasal (Valtoco®) is indicated for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from usual seizure pattern in epilepsy participants 6 years of age or older. Lastly, midazolam intranasal (Nayzilam®) is indicated for acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from usual seizure pattern in epilepsy participants 12 years of age or older. All three agents carry the same contraindication of acute narrow-angle glaucoma. Clinical guidelines from both the American Epilepsy Society and Neurocritical Care Society recommend diazepam rectal and midazolam intranasal when parenteral benzodiazepines are not available. These guidelines were also published prior to the approval of diazepam intranasal therefore, this agent lacks guideline recommendations.

Total program savings for the PDL classes will be regularly reviewed.

Program-Specific Information:	Preferred Agents	Non-Preferred Agents
	<ul style="list-style-type: none"> • Diastat® • Diazepam Rectal • Nayzilam® • Valtoco® 	<ul style="list-style-type: none"> • N/A

Type of Criteria: Increased risk of ADE Preferred Drug List
 Appropriate Indications Clinical Edit

Data Sources: Only Administrative Databases Databases + Prescriber-Supplied

Setting & Population

- Drug class for review: Anticonvulsants, Rescue Agents
- Age range: All appropriate MO HealthNet participants

Approval Criteria

- Documented diagnosis of seizure disorder **AND**
- ~~Failure to achieve desired therapeutic outcomes with trial on 2 or more preferred agents~~
 - ~~Documented trial period for preferred agents OR~~
 - ~~Documented ADE/ADR to preferred agents~~

Denial Criteria

- Lack of adequate trial on required preferred agents
- Therapy will be denied if all approval criteria are not met

Required Documentation

Laboratory Results:
MedWatch Form:

Progress Notes:
Other:

Disposition of Edit

Denial: Exception Code "0160" (Preferred Drug List)
Rule Type: PDL

Default Approval Period

1 year

References

- Evidence-Based Medicine Analysis: "Anticonvulsant Rescue Medications", UMKC-DIC; August 2022.
- Evidence-Based Medicine and Fiscal Analysis: "Anticonvulsants: Rescue Agents Therapeutic Class Review"-, Conduent Business Services, L.L.C., Richmond, VA; July 2021.
- Valtoco [package insert]. San Diego, CA: Neurelis, Inc.; February 2022.
- Diastat [package insert]. Bridgewater, NJ: Valeant Pharmaceuticals North America LLC; March 2021.
- Nayzilam [package insert]. Smyrna, GA: UCB, Inc.; February 2021.
- Brophy GM, Bell R, Claassen J; Neurocritical Care Society Status Epilepticus Guideline Writing Committee. Guidelines for the evaluation and management of status epilepticus. *Neurocrit Care*. 2012;17(1):3-23. doi: 10.1007/s12028-012-9695-z.
- Glauser T, Shinnar S, Gloss D, et al. Evidence-based guideline: treatment of convulsive status epilepticus in children and adults: report of the Guideline Committee of the American Epilepsy Society. *Epilepsy Curr*. 2016;16(1):48-61. doi: 10.5698/1535-7597-16.1.48
- Holsti M, Dudley N, Schunk J, et al. Intranasal midazolam vs rectal diazepam for the home treatment of acute seizures in pediatric patients with epilepsy. *Arch Pediatr Adolesc Med*. 2010;164(8):747-756. doi: 10.1001/archpediatrics.2010.130
- Bhattacharyya M, Kalra V, Gulati S. Intranasal midazolam vs rectal diazepam in acute childhood seizures. *Pediatr Neurol*. 2006;34(5):355-359. doi: 10.1016/j.pediatrneurol.2005.09.006

SmartPA PDL Proposal Form

© 2023 Conduent Business Services, LLC. All rights reserved. Conduent™ and Conduent Design™ are trademarks of Conduent Business Services, LLC in the United States and/or other countries.

Other company trademarks are also acknowledged.

- Fişgin T, Gurer Y, Teziç T, et al. Effects of intranasal midazolam and rectal diazepam on acute convulsions in children: prospective randomized study. *J Child Neurol.* 2002;17(2):123-126. doi: 10.1177/08830738020170020
- USPDI, Micromedex; 2022.
- Facts and Comparisons eAnswers (online); 2022 Clinical Drug Information, LLC.